

On smooth moduli space of Riemann surfaces

Yong Hou*

Institute for Advanced Study

Abstract

In this paper we study the smooth moduli space of closed Riemann surfaces. This smooth moduli is an infinite cover of the usual moduli space $\mathcal{M}_g$ of closed Riemann surfaces, and is identified with the Schottky space of rank g . The main theorem of the paper is: Closed Riemann surfaces are uniformizable by Schottky groups of Hausdorff dimension less than one. This work seem to be the only paper in literature to study question of Riemann surface uniformization and its Hausdorff dimension. We develop new techniques of rational norm of homological marking of Riemann surface and, decomposition of probability measures to prove our result. As an application of our theorem we have existence of period matrix of Riemann surface in coordinates of smooth moduli space.

Contents

1	Introduction and Main Theorem	2
2	Covering of Moduli space $\mathcal{M}_g$	5
3	Probability Measures and $\ \mathbb{W}_\Gamma\ _x$ on Schottky Space	6
4	Hausdorff Dimension of Handlebody	10

*Supported by Ambore Mondell Fundation

5	Rational Norm Q of Homological Markings on R_g and extremal length	16
6	Pants decomposition and bound of $Q([c])$	20
7	Proof of Theorem 1.1	29

1 Introduction and Main Theorem

The main theorem of this paper is:

Theorem 1.1. *Every closed Riemann surface can be uniformized by a Schottky group of Hausdorff dimension < 1 .*

Throughout this paper R_g denotes closed Riemann surface of genus g . A Kleinian group Γ is, finitely generated and discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$. Denote by Λ_Γ , the limit set of Γ , which is minimal closed Γ -invariant, nowhere dense, non-discrete, perfect subset of $\overline{\mathbb{C}}$. We say Γ is of Hausdorff dimension $\mathfrak{D}_\Gamma$ if Λ_Γ have Hausdorff dimension of $\mathfrak{D}_\Gamma$. The region of discontinuity of Γ is $\Omega_\Gamma = \overline{\mathbb{C}} - \Lambda_\Gamma$. By Ahlfors theorem, Ω_Γ/Γ is of finite many components and each is of analytically finite type Riemann surface.

Rank- g Schottky group Γ is, g -generated free, purely loxodromic Kleinian group. Equivalently, Γ is Schottky group if it is convex-cocompact representation of rank- g free group $\mathbb{F}_g$ into $\mathrm{PSL}(2, \mathbb{C})$. In particular, $\mathbb{H}^3/\Gamma$ is a handle-body and Ω_Γ/Γ is a Riemann surface.

Uniformization of R_g by Schottky group Γ , see next section for details is, $R_g = \Omega_\Gamma/\Gamma$. Theorem 1.1 states that: there exists a Schottky group Γ such that: $R_g = \Omega_\Gamma/\Gamma$ and $\mathfrak{D}_\Gamma < 1$.

The main difficulty of Theorem 1.1 is the lack of relation of uniformization Γ and its Hausdorff dimension. In fact, the question of what possible Hausdorff dimensions of Kleinian groups can uniformize R_g has not been studied at all before.

There are some interesting consequences of Theorem 1.1, which we will address separately in subsequent works. In this paper, we will state at the end an immediate simple consequence of Theorem 1.1 which implies existence of period matrix in smooth moduli coordinates.

Strategy of the proof of Theorem 1.1

First we derive a criteria for handle-body to be uniformizable by a classical Schottky group. To do so, we establish an new relation between Hausdorff dimension of limit set of Γ with it's primitive elements. This is done by a generalization of paradoxically decomposing [9, 17] the probability measure on the limit set to its generators and integrate over all primitive bases. From this decomposition we derive an new family of probability measures supported locally about each generators. By using relations among this family of probability measures, we derive an lower bound on the growth of the *mean primitive displacement* of generating elements of Γ . We show that the mean primitive displacement growth implies upper bounds on Hausdorff dimension. The measure decomposition is done abstractly on Cayley graph of free group in Section 3. The growth estimate is done for handle-body and is given in Section 4.

Secondly, that given a Riemann surface R_g , it is identified as the conformal boundary of infinity $\partial_\infty M$ of hyperbolic 3-manifold $M = (\mathbb{H}^3 \cup \Omega_\Gamma)/\Gamma$. This conformal identification is through markings on R_g . More precisely, It is classical known fact that [3, 5], we have a morphism ϕ from canonical basis with orientation of $H_1(R_g, \mathbb{Z})$ into Schottky space. This is done by representing half basis of a given basis by conformal maps of $\overline{\mathbb{C}}$. Every half basis of a given canonical basis generates a subgroup of $H_1(R_g, \mathbb{Z})$. The morphism establishes a injective corresponds of these subgroups into the Schottky space. We call a given half basis of a canonical basis a homological marking. However, it is completely unknown when a given point of Schottky space can actually cover a point of moduli space $\mathcal{M}_g$.

We study homological markings by defining a real function Q on markings of R_g . We call it *rational norm* of a marking. This is defined through ratio of geodesic representative curves with the dual geodesic to the marking geodesic as the unique minimal curve. We also define a conformal invariant of R_g , the Q -spectrum of R_g as the, collection of Q values under variation of all markings of R_g .

We then establish an inequality Lemma 5.5, between translation length of primitive elements of the image Schottky group to Q . This is done through the use of theory of extremal length. Where we use interpolation of hyperbolic metrics on planar domains.

Next we show that there exists some marking such that Q is bounded below by $\frac{2\lambda_g}{\pi}g \log(2g)$ for some $\lambda_g > 2$. We call a marking such that Q satis-

fies this lower bound is *positive*. This is done by study geodesic length ratios through elementary arcs, which are part of the marking curve on pants decompositions of R_g . The existence proof of Lemma 6.1 is by contradiction. Under assumption, we show by computation, one can always choose elementary arcs to construct curves for $[c]$ so that $Q([c])$ achieve value greater than $\frac{2\lambda_g}{\pi}g \log(2g)$. These computations are done on pair of pants.

Next we show that there exists some marking such that Q satisfies the lower bound inequality as in Lemma 5.5 under action of $\text{stab}(\phi) \subset Sp(2g, \mathbb{Z})$, stabilizer subgroup of ϕ . This is Proposition 6.4. The idea is that, by taken a marking provided by Lemma 6.1, we can show either this marking stays positive under elementary matrices E_{lm} of $\text{stab}(\phi)$ or, there must exists another marking that have larger value of Q which will be positive under E_{lm} . In fact, the ratio of lengths will increase under Dehn twist.

Proposition 6.4 implies that there exists a marking such that all primitive elements of the Schottky group will have the desired mean displacement on the handle-body. This is show in Proposition 6.6.

Finally we use these marking estimates and result on handle-body Hausdorff dimension estimates to finish proof of the main theorem. As an immediate application of Theorem 1.1, we can explicitly express period matrix of points of $\mathcal{M}_g$ in coordinates of Schottky space.

Acknowledgement

This work is made possible by unwavering supports and brilliantly insightful conversations from Peter Sarnak, whom the author is greatly indebted to. I wish to express my deep gratitude and sincere appreciation to Dave Gabai for continuous amazing support and many fundamentally influential conversations which greatly impacted my works. I wish to express my deep gratitude to Ian Agol for many enlightening and brilliant brainstorming conversations. I also want to express my greatest appreciation to Benson Farb for been able to support and help me mathematically under any circumstance without wavering. These people are critical to me to make this work possible.

I especially want to express my greatest appreciation to Ian Agol for spending very long time with me to go through the details of the computations of the paper during my visit to MSRI at Berkeley. We corrected many typos in the previous draft and also a correction to a statement. Our discussion of this work over the days at MSRI are incredibly valuable.

This paper is dedicate to: Ying Zhou and her mom Junlan Wang.

2 Covering of Moduli space $\mathcal{M}_g$

We give here a very brief and basic introduction of $\mathfrak{J}_g$ as smooth moduli of a closed Riemann surface R_g . See [3], [15], [13], [25],[24] for backgrounds materials and some of the details.

Schottky group Γ of rank g is defined as convex-cocompact discrete faithful representation of $\mathbb{F}_g$ in $\text{PSL}(2, \mathbb{C})$. It follows that Γ is freely generated by purely loxodromics $\{\gamma_i\}_1^g$. This implies we can find collection of disjointed closed topological disks $D_i, D'_i, 1 \leq i \leq g$ in the Riemann sphere $\partial\mathbb{H}^3 = \overline{\mathbb{C}}$ with boundary curves $\partial D_i = \Delta_i, \partial D'_i = \Delta'_i$. By definition Δ_i are closed Jordan curves in Riemann sphere $\partial\mathbb{H}^3$, such that $\gamma_i(\Delta_i) = \Delta'_i$ and $\gamma_i(D_i) \cap D'_i = \emptyset$. Whenever there exists a set $\{\gamma_i, \dots, \gamma_g\}$ of generators with all Δ_i, Δ'_i as circles, then it is called a classical Schottky group with $\{\gamma_1, \dots, \gamma_g\}$ classical generators.

Schottky space $\mathfrak{J}_g$ is defined as space of all rank g Schottky groups up to conjugacy by $\text{PSL}(2, \mathbb{C})$. By normalization, we can chart $\mathfrak{J}_g$ by $3g-3$ complex parameters. Hence $\mathfrak{J}_g$ is $3g-3$ dimensional complex manifold. The biholomorphic $\text{Auto}(\mathfrak{J}_g)$ group is $\text{Out}(\mathbb{F}_g)$, which is isomorphic to subgroup of the handle-body group. On the other hand, $\mathfrak{J}_{g,o}$ is not submanifold. In fact, it is nontrivial result due to Marden that $\mathfrak{J}_{g,o}$ is non-empty and non-dense set of $\mathfrak{J}$. However, it follows from a theorem of Hou [16], $\mathfrak{J}_{g,o}^\lambda$ is $3g-3$ dimensional complex submanifold. Here $\mathfrak{J}_{g,o}^\lambda$ denotes space of classical Schottky groups of Hausdorff dimension $< \lambda$.

Let $\mathcal{T}_g$ and Mod_g be the Techimuller space of R_g and it's mapping class group respectively. $\mathcal{T}_g$ is the universal cover of $\mathfrak{J}_g$. In fact, there exists subgroup $\text{Mod}_g^\phi \subset \text{Mod}_g$ which depends on a given symplectic-morphism $\phi : \pi_g \rightarrow \mathbb{F}_g$. The dependence of ϕ is only up to conjugacy within Mod_g . It follows Mod_g^ϕ is *infinite index*, torsion-free subgroup of Mod_g . Since $\mathcal{T}_g/\text{Mod}_g = \mathcal{M}_g$, we have the following commutative diagram of holomorphic covers:

$$\begin{array}{ccc} \mathcal{T}_g & \xrightarrow{\pi_U} & \mathfrak{J}_g \\ & \searrow \pi_F \quad \swarrow \pi_S & \\ & \mathcal{M}_g & \end{array}$$

In particular, $\mathfrak{J}_g$ is infinite cover of $\mathcal{M}_g$, hence can be considered *smooth* moduli of R_g . We end this section by restating Theorem 1.1:

Theorem 2.1. *Let $[R_g] \in \mathcal{M}_g$, there exist $\Gamma \in \pi_S^{-1}([R_g])$ of $\mathfrak{D}_\Gamma < 1$.*

3 Probability Measures and $\|\mathbb{W}_\Gamma\|_x$ on Schottky Space

Let Γ be a finitely generated free group with generating set ω . We denote the collection of all generating sets by W_Γ . For Schottky group with chosen point x of hyperbolic space $\mathbb{H}^3$, we also denote the minimal generator of ω with respect to x by $\mathfrak{w}_x(\omega) = \sum_{\alpha \in \omega} \frac{\text{dist}(\alpha x, x)}{g}$.

We also denote the Cayley graph of Γ with symmetric generating set $S_\omega = \omega \cup \omega^{-1}$ by CT_{S_ω} .

Definition 3.1. For Γ Schottky group, we define:

- $\mathbb{W}_\Gamma$ the set of collection of all free basis ω of Γ ,
- $\|\mathbb{W}_\Gamma\|_x$ of Γ at point x as $\|\mathbb{W}_\Gamma\|_x = \inf_{\omega \in \mathbb{W}_\Gamma} \mathfrak{w}_x(\omega)$.

We call $\|\mathbb{W}_\Gamma\|_x$, the *mean norm* of $\mathbb{W}_\Gamma$.

Theorem 3.2. *Let Γ be a Schottky group of rank g and Hausdorff dimension $\mathfrak{D}_\Gamma$. There exist nonatomic Borel measure σ_x on $\Lambda_\Gamma \times \mathbb{W}_\Gamma$ of total mass < 1 such that,*

$$\mathfrak{D}_\Gamma \leq \frac{\log\left(\frac{g}{\sigma_x(\Lambda_\Gamma \times \mathbb{W}_\Gamma)}\right)}{\|\mathbb{W}_\Gamma\|_x}.$$

Let Γ be Schottky group. Suppose $\omega_m = \langle \gamma_1, \dots, \gamma_g \rangle$ is the basis of minimal translational length T_{γ_j} . Then we can estimate $\sigma_x(\Lambda_\Gamma \times \mathbb{W}_\Gamma)$ based on relations among elements of this minimal basis.

Corollary 3.3. *Let Γ be a Schottky group. Suppose there exists $\lambda > 0$ such that $T_{\gamma_j} \leq \lambda T_{\gamma_i}$ for $\gamma_i, \gamma_j \in \omega_m$. Then there exists $x \in \mathbb{H}^3$ such that,*

$$\mathfrak{D}_\Gamma \leq \frac{(\lambda - 1) \log(2) + (\lambda + 1) \log(g)}{\|\mathbb{W}_\Gamma\|_x}.$$

Theorem 3.2 is proved by constructing families of Borel probability measures on the limit set $\Lambda(\Gamma)$. These probability measures are averages over W_Γ of measures associated to elements of W_Γ .

We first state a decomposition lemma on $(CT_S, \mathscr{W})$, with respect to some chosen symmetric generating set S_ω and word metric $\mathscr{W}$. Note that $(CT_S, \mathscr{W})$ is Gromov hyperbolic, or δ -hyperbolic.

Let μ_o be quasi-conformal measure, which is the Patterson-Sullivan measure [8], on $(CT_S, \mathcal{W})$. It is given by:

$$c_\Gamma^{-1} \Psi_o^{\mathfrak{D}_\Gamma}(\gamma^{-1}o, \zeta) \leq \frac{d\gamma^* \mu_o}{d\mu_o}(\zeta) \leq c_\Gamma \Psi_o^{\mathfrak{D}_\Gamma}(\gamma^{-1}o, \zeta), \quad c_\Gamma > 0.$$

Where $\Psi_o^{\mathfrak{D}_\Gamma}(\gamma^{-1}o, \zeta) = e^{-\mathfrak{D}_\Gamma B_o(\gamma^{-1}o, \zeta)}$, with $B_o(\gamma^{-1}o, \zeta)$ the Buseman function. We normalize so that μ_o is always taken to be probability measure on ∂CT . In addition, when we choose a indexing of $\mathbb{W}_\Gamma$, we denote it by ω_j .

Lemma 3.4. *There exists Borel measures σ_o^+, σ_o^- on $\partial CT \times \mathbb{W}_\Gamma$ with σ_o^+ a probability measure and, for each $\omega \in \mathbb{W}_\Gamma$ we have a family of Borel measures $\{\nu_{o,\gamma}\}_{\gamma \in \omega}$ and $\{\nu_{o,\gamma^{-1}}\}_{\gamma \in \omega}$ on ∂CT and $C_\omega \in (0, c_\Gamma), c_\omega \in (c_\Gamma^{-1}, c_\Gamma)$ of the following properties:*

•

$$\eta_{o,\omega}^+ = \sum_{\gamma \in \omega} \nu_{o,\gamma}, \eta_{o,\omega}^- = \sum_{\gamma \in \omega} \nu_{o,\gamma^{-1}},$$

$\eta_{o,\omega}^+$ is probability measures on ∂CT and $\eta_{o,\omega}^-(\partial CT) < 1$, for each $\omega \in \mathbb{W}_\Gamma$.

•

$$\frac{d\sigma_o^+}{d\sigma_o^-}(\zeta, \omega) = \frac{d\eta_{o,\omega}^+}{d\eta_{o,\omega}^-}(\zeta).$$

•

$$\frac{1}{(g-1) + C_\omega} \int_{\partial CT} c_\omega \Psi_o^{\mathfrak{D}_\Gamma}(\gamma^{-1}o, \zeta) d\nu_{o,\gamma^{-1}} = \int_{\partial CT} d\nu_{o,\gamma}; \text{ for } \gamma \in \omega.$$

The proof of next lemma will be based on a generalization of Culler-Shalen paradoxical measure decompositions for isometry groups of $\mathbb{H}^3$ [17]. These paradoxical decomposition were first done by Culler-Shalen [9] for the case of 2-generated cocompact Kleinian group, and later generalized in [17] to all free finitely generated subgroups of $\text{PSL}(2, \mathbb{C})$. They were interested in the embedded Margulis tube volume bounds.

Lemma 3.5. *There exists collection of Borel measures $\{\nu_\gamma\}_{\gamma \in \omega}, \{\nu_{o,\gamma^{-1}}\}_{\gamma \in \omega}$ on ∂CT such that, $\sum_{\gamma \in \omega} \nu_{o,\gamma} = \eta_{o,\omega}^+$ is probability measure and, $\sum_{\gamma \in \omega} \nu_{o,\gamma^{-1}} =$*

$\eta_{o,\omega}^-$ is measures of total mass < 1 . In addition, there exists $c_\omega \in (c_\Gamma^{-1}, c_\Gamma)$ such that,

$$\sum_{\gamma \in \omega} \int_{\partial C\Gamma} c_\omega \Psi_o^{\mathfrak{D}\Gamma}(\gamma^{-1}o, \zeta) d\nu_{o,\gamma^{-1}} = (g-1) + C_\omega.$$

Where $C_\omega \in (0, c_\Gamma)$. In addition we have,

$$\frac{1}{(g-1) + C_\omega} \int_{\partial C\Gamma} c_\omega \Psi_o^{\mathfrak{D}\Gamma}(\gamma^{-1}o, \zeta) d\nu_{o,\gamma^{-1}} = \int_{\partial C\Gamma} d\nu_{o,\gamma}.$$

Proof. Let $\mathcal{S} = \omega \cup \omega^{-1}$. Let us write every element $\gamma \in \Gamma$ as a reduced word $w_1 \cdots w_n$ with $\{w_j\} \subset \mathcal{S}$. Then we have the decomposition of Γ as $\Gamma = \{1\} \coprod \coprod_{\gamma \in \mathcal{S}} I_\gamma$, where I_γ is the set of nontrivial elements in Γ with initial letter γ . By the fact that Γ act freely on $C\Gamma$ we have $C\Gamma = \Gamma o = \{o\} \coprod \coprod_{\gamma \in \mathcal{S}} V_\gamma$ where $V_\gamma = \{wo : w \in I_\gamma\}$. Let $\mathcal{V}$ denote the collection consisting of all sets of the form $\coprod_{\gamma \in \mathcal{S}'} V_\gamma$ or $\{o\} \coprod \coprod_{\gamma \in \mathcal{S}'} V_\gamma$ for $\mathcal{S}' \subset \mathcal{S}$.

By Poincare series delta-mass construction,

$$\lim_{s \searrow \mathfrak{D}} \frac{\sum_{v \in V_\gamma} \exp(-s \text{dist}(o, v)) \delta_v}{\sum_{v \in V_\gamma} \exp(-s \text{dist}(o, v))},$$

with $C\Gamma$ and V_γ , [17] Proposition 2.5, we get a family of Borel measures $(\mu_{y,V_\gamma})_{y \in C\Gamma}$ for each $\gamma \in \mathcal{S}$, and $\text{supp}(\mu_{y,V_\gamma}) = \bar{V}_\gamma \cap \partial C\Gamma$. In general, we need to p Then, $\mu_{o,C\Gamma}$ is a probability measure on $\partial C\Gamma$. In fact, $\mu_{o,C\Gamma}$ is the normalized Patterson-Sullivan measure centered at point o . Define $\rho_\gamma := \mu_{o,V_\gamma}$ for each $\gamma \in \mathcal{S}$. By the above decomposition of $C\Gamma$, we have $\mu_{o,C\Gamma} = \mu_{o,o} + \sum_{\gamma \in \mathcal{S}} \rho_\gamma$. But $\mu_{o,o} = 0$.

Since $C\Gamma = V_{\gamma^{-1}} \coprod \gamma^{-1}V_\gamma$, we have $\gamma^{-1}V_\gamma \in \mathcal{V}$. Then by quasiconformal transformation property of Patterson-Sullivan measure, we get

$$\mu_{\gamma o, V_\gamma} = \gamma^{-1*} \mu_{o, C\Gamma - V_{\gamma^{-1}}} = \gamma^{-1*} (\mu_o - \rho_{\gamma^{-1}}).$$

This implies we have,

$$d\mu_{\gamma o, V_\gamma} = c_\gamma \Psi_o^{\mathfrak{D}\Gamma}(\gamma^{-1}o, \xi) d\mu_{o, V_\gamma}, \text{ for some } c_\gamma \in (c_\Gamma^{-1}, c_\Gamma).$$

From this, we get

$$\int_{\partial C\Gamma} c_\gamma \Psi_o^{\mathfrak{D}\Gamma}(\gamma^{-1}o, \xi) d\rho_{\gamma^{-1}} = \int_{\partial C\Gamma} d(\gamma^{-1*}(\mu_{o,C\Gamma} - \rho_\gamma)) = 1 - \int_{\partial C\Gamma} d\rho_\gamma.$$

Since $\mu_{o,CT} = \sum_{\gamma \in \omega} \rho_\gamma + \sum_{\gamma \in \omega} \rho_{\gamma^{-1}}$, we have one of $\sum_{\gamma \in \omega} \rho_\gamma, \sum_{\gamma \in \omega} \rho_{\gamma^{-1}}$ must have total weight over ∂CT of $\geq \frac{1}{2}$. This is one of the property two that we exploit later in next corollary to show lower bounds of σ_o^- based on indexing choice of $\mathbb{W}_\Gamma$.

Set $C_\omega = \sum_{\gamma \in \omega} \int_{\partial CT} d\rho_{o,\gamma^{-1}}$. There exists $c_\omega \in (c_\Gamma^{-1}, c_\Gamma)$ such that,

$$\nu_{o,\gamma^{-1}} = \rho_{o,\gamma^{-1}},$$

$$\nu_{o,\gamma} = \frac{\mu_o - \rho_{o,\gamma}}{c_\omega(g-1) + c_\omega \sum_{\gamma \in \omega} \int_{\partial CT} d\rho_{o,\gamma^{-1}}},$$

satisfies our conditions. $\square$

Proof. Proposition 3.4:

For $\omega \in \mathbb{W}_\Gamma$, define measures by $\eta_{o,\omega}^+ = \sum_{\gamma \in \omega} \nu_{o,\gamma}$, and $\eta_{o,\omega}^- = \sum_{\gamma \in \omega} \nu_{o,\gamma^{-1}}$. By Lemma 3.5, $\eta_{o,\omega}^+$ is a probability measures on ∂CT .

Index $\mathbb{W}_\Gamma = \cup_{i \geq 1} \omega_i$. Set $\sigma_{o,N}^+ = \frac{1}{N} \sum_{i=1}^N \eta_{o,\omega_i}^+ \otimes \delta_{\omega_i}$, $\sigma_{o,N}^- = \frac{1}{N} \sum_{i=1}^N \eta_{o,\omega_i}^- \otimes \delta_{\omega_i}$. We extend $\sigma_{o,N}^+, \sigma_{o,N}^-$ to $\partial CT \times \mathbb{W}_\Gamma$ trivially by setting it to be zero on $\mathbb{W}_\Gamma - \cup_{j \leq N} \omega_j$. Note that since η_{o,ω_j}^+ are probability measures, we have for all N ,

$$\int_{\partial CT \times \mathbb{W}_\Gamma} d\sigma_{o,N}^+ = 1.$$

Since $\eta_{o,\omega_j}^-(\partial CT) < 1$ for all j we have,

$$\int_{\partial CT} d\sigma_{o,N}^- < 1.$$

We have the weak-limit of, $\sigma_n^+ \rightarrow \sigma_o^+$ and $\sigma_n^- \rightarrow \sigma_o^-$. By construction we have $\sigma_o^\pm$ satisfies our conditions. $\square$

Corollary 3.6. *There exists a sequence of Borel measures $\sigma_{o,n}^\pm$ on $\partial CT \times \mathbb{W}_\Gamma$ such that, support of $\sigma_{o,n}^\pm$ is $\partial CT \times \cup_{j \leq n} \omega_j$ for a given index of $\mathbb{W}_\Gamma$ and, $\sigma_{o,n}^\pm \rightarrow \sigma_o^\pm$ weakly. In addition, σ_o^+ is a probability measure, so it's total weight is independent on index of $\mathbb{W}_\Gamma$, and there exists a indexing of $\mathbb{W}_\Gamma$ such that, $\sigma_o^-(\partial CT \times \mathbb{W}_\Gamma) \geq \frac{1}{2}$.*

Proof. Since $\eta_{o,n}^+$ for all n are probability measures, we can easily see by our construction in the proof of Proposition 3.4 that $\sigma_{o,n}^+ \rightarrow \sigma_o^+$ is indexing independent of $\mathbb{W}_\Gamma$.

We can choose a indexing of $\mathbb{W}_\Gamma$ such that given ω_i we have either $\omega_{i+1}, \omega_{i-1}$ is the inverse of ω_i . Since $\mu_o = \eta_{o, \omega_i}^- + \eta_{o, \omega_i^{-1}}^-$ we have by using this indexing, $\sigma_{o, n}^-(\partial CT \times \mathbb{W}_\Gamma) \rightarrow \frac{\mu_o}{2}(\partial CT)$. Alternatively, if given a indexing such that $\sigma_o^-(\partial CT \times \mathbb{W}_\Gamma) < \frac{1}{2}$ then we can replace all ω_i with ω_i^{-1} . This would imply $\frac{1}{N} \sum_{i=1}^N \eta_{o, \omega_i}^-(\partial CT \times \mathbb{W}_\Gamma) \geq \frac{1}{2}$ for large N . Hence we have required result. $\square$

4 Hausdorff Dimension of Handlebody

Take Γ to Schottky group. Then the parameter in Lemma 3.5 is $c = 1$.

Let Λ_Γ be the limit set of Γ , which is the minimal invariant uniformly perfect closed no-where dense subset of $\mathbb{C}$. The open invariant set $\Omega_\Gamma = \mathbb{C} - \Lambda_\Gamma$ is the region of discontinuity of Γ . Then $\mathfrak{D}_\Gamma$ from previous section is the Hausdorff dimension of Λ_Γ , which we denote by $\mathfrak{D}_\Gamma$. We also have ∂CT is identified with Λ_Γ . Since Γ is convex-cocompact, CT is quasi-isometric to CH_Γ , convex hull of Γ . In addition, the density $\Psi_x(\gamma^{-1}x, \xi)$ is the Poisson kernel on $\mathbb{H}^3 \cup \overline{\mathbb{C}}$. Explicitly we have [17],

$$\Psi_x(\gamma^{-1}x, \xi)^{\mathfrak{D}_\Gamma} = \left(\frac{1}{\cosh(\text{dist}(x, \gamma x)) - \sinh(\text{dist}(x, \gamma x)) \cos \angle \gamma^{-1}xx\xi} \right)^{\mathfrak{D}_\Gamma}.$$

Here $\text{dist}(x, y)$ is the hyperbolic distance for $x, y \in \mathbb{H}^3$. Hence we can proceed explicit computation of measure over Λ_Γ .

We summary this in the following:

Corollary 4.1. *Let Γ be Schottky group. Lemma 3.5 holds for $c = 1$ and $\Psi_o^{-\mathfrak{D}_\Gamma}(\gamma o, \zeta)$ given by the Poisson Kernel.*

We denote $\Psi_x(\gamma^{-1}x, 0)$ for $\Psi_x(\gamma^{-1}x, \xi)$ when $\angle \gamma^{-1}xx\xi = 0$. This is simply $\exp(\text{dist}(x, \gamma x))$.

Given $\alpha \in \omega$ we define, $\omega(\alpha)$ is the generating set given by α and $\{\gamma\alpha\}_{\gamma \in \omega - \alpha}$. This is the generating set given by *shifting* the original ω by α .

Next we will prove a lemma that will bound the mean displacement of any given ω at a point to that of the derivative of $\eta_{x, \omega(\alpha)}^-$ with respect to some transformed measure. This will provide connection between the mean displacement of ω with measures $\sigma_x^\pm$.

Lemma 4.2 (Shifting lemma). *Define $d\eta_{x,\omega}^*(\zeta) = \sum_{\gamma \in \omega} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) d\nu_{x,\gamma^{-1}}$. Then there exists $\alpha \in \omega$ such that,*

$$\frac{d\eta_{x,\omega}^*}{d\eta_{x,\omega(\alpha)}^-}(\zeta) \geq \sum_{\gamma \in \omega} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g}.$$

Remark 4.3. Note that $\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)$ is notation for $\exp(\mathfrak{D}_\Gamma \text{dist}(\gamma^{-1}x, x))$.

Recall that Patterson-Sullivan measures provides exact quasiconformal distortions under group transformation. The idea of this lemma is that, we want to somehow gauge the distortion created when we change the measure $\nu_{x,\gamma}$ under transformations. More precisely, we want to estimate the distortion under Nielsen transformation. The point of shifting lemma is to estimate the lower bound of this distortion, which states that the distortion is at least the average of overall distortion.

Proof. Let $\delta > 0$ small. Define subset $E_\delta(x, \gamma) = \{\zeta \in \Lambda_\Gamma | \angle \gamma^{-1}x\zeta < \delta\}$. Let $\alpha_\delta \in \omega$ such that

$$\nu_{x,\alpha_\delta^{-1}}(E_\delta(x, \alpha_\delta)) \leq \nu_{x,\gamma^{-1}}(E_\delta(x, \gamma)), \text{ for all } \gamma \in \omega.$$

Note that $\text{supp}(\nu_{x,\gamma^{-1}}) = \bar{V}_{\gamma^{-1}} \cap \Lambda_\Gamma$. For $\beta \in \omega(\alpha_\delta) - \alpha_\delta$ we have,

$$\text{supp}(\nu_{x,\beta^{-1}}) = \text{supp}(\nu_{x,\alpha_\delta^{-1}\gamma^{-1}}) \subset \text{supp}(\nu_{x,\alpha_\delta^{-1}}),$$

$$\nu_{x,\beta^{-1}} \leq \nu_{x,\alpha_\delta^{-1}}.$$

This implies that for $\gamma' \in I_{\alpha_\delta^{-1}}$ (word start with α_δ) we have,

$$\nu_{x,\alpha_\delta^{-1}}(E_\delta(x, \gamma')) \geq \nu_{x,\beta^{-1}}(E_\delta(x, \gamma')), \text{ for all } \beta \in \omega(\alpha_\delta).$$

Hence we have,

$$\begin{aligned}
& \lim_{\delta \rightarrow 0} \frac{\sum_{\gamma \in \omega} \int_{E_\delta(x, \gamma')} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) d\nu_{x, \gamma^{-1}}}{\sum_{\gamma \in \omega(\alpha_\delta)} \int_{E_\delta(x, \gamma')} d\nu_{x, \gamma^{-1}}} \\
& \geq \lim_{\delta \rightarrow 0} \frac{\sum_{\gamma \in \omega} \inf_{\zeta \in E_\delta(x, \gamma')} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) \int_{E_\delta(x, \gamma')} d\nu_{x, \gamma^{-1}}}{\sum_{\gamma \in \omega(\alpha_\delta)} \int_{E_\delta(x, \gamma')} d\nu_{x, \gamma^{-1}}} \\
& \geq \lim_{\delta \rightarrow 0} \frac{\sum_{\gamma \in \omega} \inf_{\zeta \in E_\delta(x, \gamma')} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) \nu_{x, \alpha_\delta^{-1}}(E_\delta(x, \gamma'))}{\sum_{\gamma \in \omega(\alpha_\delta)} \int_{E_\delta(x, \gamma')} d\nu_{x, \gamma^{-1}}} \\
& \geq \lim_{\delta \rightarrow 0} \frac{\sum_{\gamma \in \omega} \inf_{\zeta \in E_\delta(x, \gamma')} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) \nu_{x, \alpha_\delta^{-1}}(E_\delta(x, \gamma'))}{g \nu_{x, \alpha_\delta^{-1}}(E_\delta(x, \gamma'))} \\
& \geq \sum_{\gamma \in \omega} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g}
\end{aligned}$$

□

Proof. Theorem 3.2:

Let $N > 0$ be a large integer. By Lemma 3.4 and Corollary 3.6, there exists $\epsilon_N > 0$ such that,

$$\begin{aligned}
\int_{\Lambda_\Gamma \times \mathbb{W}_\Gamma} d\sigma_x^+ & \geq (1 - \epsilon_N) \sum_{1 \leq j \leq N} \frac{1}{N} \int_{\Lambda_\Gamma} d\eta_{x, \omega_j}^+ \\
& \geq (1 - \epsilon_N) \sum_{1 \leq j \leq N} \sum_{\gamma \in \omega_j} \frac{1}{N} \int_{\Lambda_\Gamma} d\nu_{x, \gamma^+} \\
& \geq (1 - \epsilon_N) \sum_{1 \leq j \leq N} \sum_{\gamma \in \omega_j} \frac{1}{N(g - 1 + C_{\omega_j})} \int_{\Lambda_\Gamma} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) d\nu_{x, \gamma^{-1}}
\end{aligned}$$

Since $c_\Gamma = 1$ and $C_{\omega_j} \in (0, 1)$ for all j we have,

$$\geq \frac{(1 - \epsilon_N)}{g} \sum_{1 \leq j \leq N} \sum_{\gamma \in \omega_j} \frac{1}{N} \int_{\Lambda_\Gamma} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) d\nu_{x, \gamma^{-1}}.$$

By shifting Lemma 4.2, for every ω_j there exists $\alpha_j \in \omega_j$ such that,

$$\frac{\sum_{\gamma \in \omega_j} \Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, \zeta) d\nu_{x, \gamma^{-1}}}{\sum_{\gamma \in \omega_j(\alpha_j)} d\nu_{x, \gamma^{-1}}} \geq \sum_{\gamma \in \omega_j} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g}.$$

Hence we have,

$$\int_{\Lambda_\Gamma \times \mathbb{W}_\Gamma} d\sigma_x^+ \geq \frac{(1 - \epsilon_N)}{g} \inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g} \sum_{1 \leq j \leq N} \sum_{\gamma \in \omega_j(\alpha_j)} \frac{1}{N} \int_{\Lambda_\Gamma} d\nu_{x, \gamma^{-1}}.$$

Note the above inequality holds for any chosen indexing of $\mathbb{W}_\Gamma$, and by Corollary 3.6, we can choose some indexing such that $\sigma_x^-(\Lambda_\Gamma \times (W)_\Gamma) \geq \frac{1}{2}$. However, our shifting is indexing dependent hence we can't in general simply just bound by $1/2$, unless we have some bounds on generators translation length.

$$\begin{aligned} \int_{\Lambda_\Gamma \times \mathbb{W}_\Gamma} d\sigma_x^+ &\geq \frac{(1 - \epsilon_N)}{g} \inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g} \sum_{1 \leq i \leq N} \sum_{\gamma \in \omega_{j_i}(\alpha_{j_i})} \frac{1}{N} \int_{\Lambda_\Gamma} d\nu_{x, \gamma^{-1}} \\ &\geq \frac{(1 - \epsilon_N)}{g} \inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}x, 0)}{g} \sum_{1 \leq i \leq N} \frac{1}{N} \int_{\Lambda_\Gamma} d\eta_{x, \omega_{j_i}(\alpha_{j_i})}^-. \end{aligned}$$

Since $\Psi_x^{\mathfrak{D}_\Gamma}(\gamma^{-1}, 0) = \exp(\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x))$,

$$\geq \frac{(1 - \epsilon'_N)}{g} \inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{e^{\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x)}}{g} \int_{\Lambda_\Gamma \times \mathbb{W}_\Gamma} d\sigma_x^-.$$

By $\sigma_x^+(\Lambda_\Gamma \times \mathbb{W}_\Gamma) = 1$ and $\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)$ we have,

$$1 \geq \frac{(1 - \epsilon'_N)\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)}{g} \inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{e^{\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x)}}{g}.$$

This implies that,

$$\inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{e^{\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x)}}{g} \leq \frac{g}{(1 - \epsilon'_N)\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)}.$$

From the inequality,

$$\inf_{\omega \in \mathbb{W}_\Gamma} \sum_{\gamma \in \omega} \frac{\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x)}{g} \leq \inf_{\omega \in \mathbb{W}_\Gamma} \log\left(\sum_{\gamma \in \omega} \frac{e^{\mathfrak{D}_\Gamma \text{dist}(x, \gamma^{-1}x)}}{g}\right),$$

and since ϵ'_N is arbitrarily small, hence we have,

$$\mathfrak{D}_\Gamma \leq \frac{\log\left(\frac{g}{\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)}\right)}{\|\mathbb{W}_\Gamma\|_x}.$$

□

To give *best* possible estimate of $\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)$ we need some control of the shifting of $\mathbb{W}_\Gamma$. This shifting information is provided by the the relationship among the generators. The idea is to pick the best possible indexing of $\mathbb{W}_\Gamma$ such that the shifting will remain sufficiently bounded below along the indexing.

Corollary 4.4. *Let Γ be a Schottky group such that there exists $x \in \mathbb{H}^3$ and indexing of $\mathbb{W}_\Gamma = \cup_j \{\omega_j\}$ such that all shifting $\omega_j(\alpha_j)$ have $\nu_{x,\gamma^{-1}} \geq \frac{1}{2g}$ for $\omega_j(\alpha_j)$ then,*

$$\mathfrak{D}_\Gamma \leq \frac{2 \log(g)}{\|\mathbb{W}_\Gamma\|_x}.$$

We can obtain an similar estimate with average bounds on $\nu_{x,\gamma^{-1}}$ as follows.

Corollary 4.5. *Let Γ be a Schottky group such that there exists $x \in \mathbb{H}^3$ and indexing of $\mathbb{W}_\Gamma = \cup_j \{\omega_j\}$ with $\nu_{x,\gamma^{-1}}(\Lambda_\Gamma) \geq \frac{1}{g-1} \sum_{\beta \in \omega_j - \gamma} \nu_{x,\beta^{-1}}(\Lambda_\Gamma)$ for every $\gamma \in \omega_j$. Then*

$$\mathfrak{D}_\Gamma \leq \frac{2 \log g}{\|\mathbb{W}_\Gamma\|_x}.$$

Let Γ be Schottky group. Suppose $\omega_m = \langle \gamma_1, \dots, \gamma_g \rangle$ is the basis of minimal translational length T_{γ_j} . Then we can estimate ν_{x,ω_j} based on relations among elements of this minimal basis.

Corollary 4.6. *Let Γ be a Schottky group. Suppose there exists $\lambda \geq 1$ such that $T_{\gamma_j} \leq \lambda T_{\gamma_i}$ for $\gamma_i, \gamma_j \in \omega_m$. Then there exists $x \in \mathbb{H}^3$ such that,*

$$\mathfrak{D}_\Gamma \leq \frac{(\lambda - 1) \log(2) + (\lambda + 1) \log(g)}{\|\mathbb{W}_\Gamma\|_x}.$$

Proof. For convex cocompact Γ , Patterson-Sullivan measure is the unique $\mathfrak{D}_\Gamma$ -Hausdorff measure. This implies $\nu_{x,g^{-1}}, \gamma \in \omega$ is absolutely continuous to the Hausdorff measure of support $\bar{V}_{g^{-1}} \cap \Lambda_\Gamma$. By $T_{\gamma_j} \leq \lambda T_{\gamma_i}$, we can choose $x \in \mathbb{H}^3$ such that $\nu_{x,\gamma_k^{-1}}(\Lambda_\Gamma) \geq \frac{1}{2g}$ for some $\gamma_k \in \omega_m$, and the total mass of $\nu_{x,\gamma_j^{-1}}$ is bounded by $\nu_{x,\gamma_j^{-1}}(\Lambda_\Gamma) \geq \nu_{x,\gamma_i^{-1}}^\lambda(\Lambda_\Gamma)$. Let $\omega_m(\gamma_j) = \{\gamma_j, \gamma_i \gamma_j\}_{i \neq j}$ be the shifted basis. Since $\sum_{\gamma \in \omega_m(\gamma_j) - \gamma_j} \nu_{x,\gamma^{-1}}(\Lambda_\Gamma) = \nu_{x,\gamma_j^{-1}}(\Lambda_\Gamma)$ we have,

$$\begin{aligned} \sum_{\gamma \in \omega_m(\gamma_j)} \nu_{x,\gamma^{-1}}(\Lambda_\Gamma) &\geq \nu_{x,\gamma_k^{-1}}^\lambda(\Lambda_\Gamma) + \sum_{\gamma \in \omega_m(\gamma_j) - \gamma_j} \nu_{x,\gamma^{-1}}(\Lambda_\Gamma) \\ &\geq \nu_{x,\gamma_k^{-1}}^\lambda(\Lambda_\Gamma) + \nu_{x,\gamma_k^{-1}}^\lambda(\Lambda_\Gamma) \geq \frac{2}{(2g)^\lambda}. \end{aligned}$$

Using the notations in the proof of Theorem 3.2, since ω_m is the minimal length generator basis, we can choose an indexing of $\mathbb{W}_\Gamma$ such that,

$$\sum_{1 \leq i \leq N} \frac{1}{N} \sum_{\gamma \in \omega_{j_i}(\alpha_{j_i})} \nu_{x, \gamma^{-1}}(\Lambda_\Gamma) \geq \frac{2}{(2g)^\lambda}.$$

This implies that, $\sigma_x(\Lambda_\Gamma \times \mathbb{W}_\Gamma) \geq \frac{2}{(2g)^\lambda}$ which give the result. $\square$

Rather than uniform bound of λ , we can relax the condition of Corollary 4.6 between T_i, T_j . Arrange $T_{\gamma_i} \leq T_{\gamma_j}$ for $i \leq j$ as follows: Suppose $T_j \leq \lambda_{ji} T_i$ for some collection of $\lambda_{ji} \geq 1, i \leq j$. We take convention that $\lambda_{ij} = 1$ for $i \leq j$. If γ_j is the shifting generator for $\omega_m(\gamma_j)$, then by replacing generator γ_k for, $k = j-1$ if $j > 1$ and $k = 2$ when $j = 1$, with its inverse in ω_m if necessary, we can assume that $\nu_{x, \gamma_k^{-1}}(\Lambda_\Gamma) \geq \frac{1}{2g}$. This implies that, $\sum_{\gamma \in \omega_m(\gamma_j)} \nu_{x, \gamma^{-1}}(\Lambda_\Gamma) \geq \frac{2}{(2g)^{\lambda'_{jk}}}$. Set $\bar{\lambda} = \sup_{jk} \lambda'_{jk}$ we have the following version of above corollary:

Corollary 4.7. *Let Γ be a Schottky group. Suppose there exists $\lambda_{ij} \geq 1$ such that $T_{\gamma_j} \leq \lambda_{ji} T_{\gamma_i}, i \leq j$ for $\gamma_i, \gamma_j \in \omega_m$. Then there exists $x \in \mathbb{H}^3$ such that,*

$$\mathfrak{D}_\Gamma \leq \frac{(\bar{\lambda} - 1) \log(2) + (\bar{\lambda} + 1) \log(g)}{\|\mathbb{W}_\Gamma\|_x}.$$

Finally, we mention couple of interesting consequences of Theorem 3.2 related to injectivity radius and classical Schottky groups.

Define $H_c = \sup\{|\lambda| \text{ such that all Schottky group of } \mathfrak{D}_\Gamma < \lambda \text{ is classical}\}$. H_c is the maximal parameter such that if Γ have Hausdorff dimension $< H_c$ then Γ is classical Schottky group. It follows from theorem of Hou [16], such that H_c exists. We have next obvious corollary:

Corollary 4.8. *There exists $\tau_c > 0$ such that any Schottky group Γ of rank g with $\|\mathbb{W}_\Gamma\|_x > \tau_c \log\left(\frac{g}{\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)}\right)$ for all $x \in \mathbb{H}^3$ is classical Schottky group.*

Proof. By a theorem of Hou [16], there exists maximal $H_c > 0$ such that any finitely generated free Kleinian Γ with Hausdorff dimension $< H_c$ is classical Schottky group. Set $\tau = \frac{1}{H_c}$ completes the proof. $\square$

Corollary 4.9. *Let $H = \mathbb{H}^3/\Gamma$ be hyperbolic handlebody of rank $g > 1$ such that $\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma) \geq 1/2$. There exists universal $\tau_c > 0$ such that, if the injectivity radius i_H of H satisfies $i_H > \tau_c \log(2g)$, then H is uniformized by classical Schottky group Γ .*

For a hyperbolic 3-manifold M , if the $\pi_1(M)$ is not free then there is imbedded surface which would put an upper bounds on the injectivity radius. Hence for sufficiently large injectivity of a given hyperbolic 3-manifold, the fundamental group must be free. So we have the following corollary:

Corollary 4.10. *Let M be a hyperbolic 3-manifold with fundamental group of rank g . If $i_M > c \log \left(\frac{g}{\sigma_x^-(\Lambda_\Gamma \times \mathbb{W}_\Gamma)} \right)$, then M is unformized by a classical Schottky group.*

5 Rational Norm Q of Homological Markings on R_g and extremal length

Let $H_1(R_g, \mathbb{Z})$ be the first homology group and denote by B_1 the set of canonical basis of $H_1(R_g, \mathbb{Z})$. This is given by class: $[\alpha_i], [\beta_i]$ satisfies $\langle [\alpha_i], [\alpha_j] \rangle = \langle [\beta_i], [\beta_j] \rangle = 0$ and $\langle [\alpha_i], [\beta_j] \rangle = \delta_{ij}$, $\langle [\beta_j], [\alpha_i] \rangle = -\delta_{ij}$.

Let $\pi : B_1 \rightarrow BH_1$ be the projection to the collection of first set of g cycles $\{\alpha_i\}_1^g$ i.e. $\langle \alpha_i, \alpha_j \rangle = 0$, half-basis. Denote B_α the subgroup of $H_1(R_g, \mathbb{Z})$ generated by $[c] \in BH_1$. We define $B_{1/2}$ to be the collection of all such subgroups B_α for $[c] \in BH_1$.

There exists a $\phi : BH_1 \rightarrow \mathfrak{J}_g$ maps into the Schottky space. The map ϕ is a morphism such that maps all $\phi(\alpha_j) = \gamma_j, 1 \leq j \leq g$. The curves $\{\alpha_1, \dots, \alpha_g\}$ are called *cut system*. Denote $\Gamma_{[c]} = \langle \gamma_1, \dots, \gamma_g \rangle$ the image Schottky group of $[c]$ under ϕ . Each B_α , subgroup of $H_1(R_g, \mathbb{Z})$ generated by $\{\alpha_1, \dots, \alpha_g\}$ uniquely determines $\Gamma_{[c]}$. Different set of α'_i which generates same subgroup of $H_1(R_g, \mathbb{Z})$ gives same Schottky group under ϕ , of different set of generators, which corresponds to α'_i cut system. We have injective map of $B_{1/2}$ into $\mathfrak{J}_g$. The fundamental domain of $\Gamma_{[c]}$ is conformally equivalent to the planar (genus zero) domain, $R_g - \cup_{1 \leq i \leq g} \alpha_i$. For details, see [3, 5].

We denote the region of discontinuity of the image Schottky group $\Gamma_{[c]}$ by $\Omega_{[c]}$. The domain $\Omega_{[c]}$ is hyperbolic planar domain with hyperbolic metric $\rho_{[c]}$, which is the $\Gamma_{[c]}$ -invariant Poincare metric of hyperbolic disk. If ρ_R is the Poincare hyperbolic metric of R_g then, we have holomorphic covering map, $\pi_s : (\Omega_{[c]}, \rho_{[c]}) \rightarrow (R_g, \rho_R)$.

B_1 is invariant under the symplectic group $Sp(2g, \mathbb{Z})$. Let $stab(\phi)$ denote the normal subgroup of stabilizer of B_α . The subgroup $stab(\phi)$ is generated by elements which corresponds to Nielsen transformations of generators of $\Gamma_{[c]}$.

Let $\theta \in Sp(2g, \mathbb{Z})$, we define the action $\theta[c]$ for $[c] \in BH_1$ as: $\pi\theta\pi^{-1}[c]$. We also set $\hat{B}H_1 = BH_1/stab(\phi)$. Denote elements of $\hat{B}H_1$ by $\llbracket c \rrbracket$.

Given $[c] = \{\alpha_i\}_1^g \in BH_1$, the collection of g cycles $\{\sigma_j\}_1^g \in \pi^{-1}[c]$ are called *dual cycles* of α_i cycles. Denote this collection by $D(\alpha)$. We also denote by $D(\alpha_i)$ to be the collection of all dual cycles to α_i in $D(\alpha)$. i.e. simple closed curves in $D(\alpha)$ of intersection 1 with α_i .

For any given curve $\sigma \subset (R_g, \rho_R)$, we denote the hyperbolic length of σ by $\ell(\sigma)$. This is also the hyperbolic length in $\pi_s^{-1}(\sigma) \subset (\Omega_{\Gamma[c]}, \rho_{[c]})$.

For $[c] \in BH_1$, Denote $[c]_i$ as α_i , the i -th cut system cycles. $\ell([c]_i)$ is the hyperbolic length of the *geodesic* representative curve of α_i . Since R_g is compact, there exists a unique $\beta_i^* \in D(\alpha_i)$, geodesic representative curve such that, $\ell(\beta_i^*) = \inf_{\beta \in D(\alpha_i)} \ell(\beta)$. We also define the following notations:

$$\|[c]_i\| = \ell^2([c]_i), \quad \|[c]_i\|_D = \ell^2(\beta_{\alpha_i}^*).$$

Remark 5.1. We make convention that for a $[c] \in BH_1$ represented by geodesics $\{\alpha_i\}_1^g$, we identify $[c]$ to a basis in $H_1(R_g, \mathbb{Z})$ by adjoining the $\beta_{\alpha_i}^*$ the unique geodesic cycles as: $\alpha_i, \beta_{\alpha_i}^*$. So we speak of $[c]$ as basis and element of BH_1 interchangeably through this identification.

Definition 5.2. Let $[c] \in BH_1$. We also define the Rational Norms $Q([c]_i)$ of $[c]$ as the collection of all $Q([c])$:

$$Q([c]_i) = \frac{\|[c]_i\|_D}{\|[c]_i\|}, \quad Q([c]) = \sum_{i=1}^g Q([c]_i).$$

Definition 5.3. We define Q_λ the Q -spectrum of R_g as:

$$Q_\lambda(R_g) = \{Q([c]) \mid [c] \in BH_1\}.$$

Note $Q_\lambda(R_g)$ is non-discrete countable set. Next proposition is obvious:

Proposition 5.4. $Q_\lambda(R_g)$ is conformal invariant and defines a set-valued function on $\mathcal{M}_g$.

Let $\{T_{[c],i}\}_1^g$ denote the collection of translation length of elements $\{\gamma_i\} \in \Gamma_{[c]}$. Set $\mathcal{T}_{[c]} = \frac{1}{g} \sum_{1 \leq i \leq g} T_{[c],i}$. We call $\mathcal{T}_{[c]}$ the *Schottky length* of $[c]$.

Given R_g a Riemann surface or domain of $\mathbb{C}$. Denote $\text{conf}(R_g)$ space of all conformal metric on R_g . Locally, $\psi \in \text{conf}(R_g)$ is given by quadratic differential $\psi(z)dz^2$.

Let Φ be a collection of curves in R_g . Recall the *extremal length* $\mathcal{E}_{R_g}(\Phi)$ is given by:

$$\mathcal{E}_{R_g}(\Phi) = \sup_{\psi \in \text{conf}(R_g)} \frac{\inf_{\sigma \in \Phi} \left(\int_{\sigma} |\psi| \right)^2}{\int_{R_g} |\psi|^2}$$

Note that it is simple fact that $\mathcal{E}_{R_g}(\Phi)$ is conformal invariant.

Next we will use extremal length to establish lower bounds of $T_{[c]_i}$ by $Q([c]_i)$.

Lemma 5.5.

$$T_{[c]_i} \geq \frac{\pi}{2} Q([c]_i).$$

Proof. Let $C_i, C'_i \subset \mathbb{C}$ be the lift of $[c]_i$ which are Jordan curves that bounds disjoint closed disks D_i, D'_i . such such: $\gamma_i(C_i) = C'_i, \gamma_i(D_i^o) \cap D_i^c$. Here γ_i is generators of $\Gamma_{[c]}$ given by $\phi([c])$. As before, $\rho_{[c]}$ is the hyperbolic metric on $\Omega_{[c]}$.

Let Φ_i be the collection of all paths connecting C_i to C'_i in $\mathbb{C}$. Let $\mathcal{R}_i = \mathbb{C} - (D_i \cup D'_i)$. Also set $\mathcal{R} = \mathbb{C} - \cup_i (D_i \cup D'_i)$, and $D_i^c = \cup_{j \neq i} (D_j \cup D'_j)$.

Note that since $\mathcal{R} \subset \Omega_{[c]}$, $\rho_{[c]}$ defines a hyperbolic metric on $\mathcal{R}$. Let w be any curve in $\mathcal{R}$ connecting D_i, D'_i . Let $\ell(w)$ be the $\rho_{[c]}$ -hyperbolic length of w .

Denote U collection of all curves in $\mathcal{R}_i$ connecting D_i, D'_i . Choose a conformal metric h_w on $\mathcal{R}_i$ such that, $\inf_{u \in U} \ell_{h_w}(u) > \ell(w)$. Let $\epsilon > 0$ such that, the D_ϵ , ϵ -neighborhood of D_i^c is: $D_\epsilon \cap (D_i \cup D'_i) = \emptyset$. Choose a $\sigma_\epsilon(z)$ smooth function of $\mathbb{C}$, which is approximate characteristic cut-off function such that:

$$\sigma_\epsilon(z) = \begin{cases} 1 & \text{if } z \in \mathcal{R} \\ 0 & \text{if } x \in D_\epsilon \end{cases}$$

and, $d\rho_{\epsilon,w}^2 = (d\rho_{[c]}^2)^{\sigma_\epsilon} (dh_w^2)^{1-\sigma_\epsilon}$ is of negatively pinched curvature. If we denote metric density by the same notation and write hyperbolic metric in conformal factors $e^{\rho_{[c]}}|dz|$ and $e^{h_w}|dz|$ we have, $\rho_{\epsilon,w} = \sigma_\epsilon \rho_{[c]} + (1 - \sigma_\epsilon)h_w$ is ϵ -family of pinched negatively curved metric on $\mathcal{R}_i$.

First we establish bounds of $Q([c]_i)$ by extremal length $\mathcal{E}_{\mathcal{R}_i}(\Phi)$.

It follows from the isoperimetric inequality for negatively pinched manifold [2, 27] we have,

$$\int_{\mathcal{R}_i} |\rho_{\epsilon,w}|^2 \leq \left(\int_{C_i \cup C'_i} |\rho_{\epsilon,w}| \right)^2.$$

$$\mathcal{E}_{\mathcal{R}_i}(\Phi_i) = \sup_{\psi \in \text{conf}(\mathcal{R}_i)} \frac{\inf_{\sigma \in \Phi_i} \left(\int_{\sigma} |\psi| \right)^2}{\int_{\mathcal{R}_i} |\psi|^2} \geq \frac{\inf_{\sigma \in \Phi_i} \left(\int_{\sigma} |\rho_{\epsilon, w}| \right)^2}{\left(\int_{\mathcal{R}_i} |\rho_{\epsilon, w}|^2 \right)}.$$

By isoperimetric inequality we have,

$$\geq \frac{\inf_{\sigma \in \Phi_i} \left(\int_{\sigma} |\rho_{\epsilon, w}| \right)^2}{\left(\int_{C_i \cup C'_i} |\rho_{\epsilon, w}| \right)^2}.$$

Since assuming ϵ is sufficiently small we have, $\int_{C_i \cup C'_i} |\rho_{\epsilon, w}| < \int_{C_i \cup C'_i} |\rho_{[c]}| + \delta_{\epsilon}$, with $\delta_{\epsilon} \rightarrow 0$. We have,

$$> \frac{\inf_{\sigma \in \Phi_i} \left(\int_{\sigma} |\rho_{\epsilon, w}| \right)^2}{\left(\int_{C_i \cup C'_i} |\rho_{[c]}| + \delta_{\epsilon} \right)^2}.$$

By $\inf_{u \in U} \ell_{h_w}(u) > \ell(w)$, implies all curves u in $\mathcal{R}_i$ connecting D_i, D'_i that intersects D_i^c must have $\ell_{\rho_{\epsilon, w}}(u) > \ell(w)$. Let V denote curves in $\mathcal{R}$ connecting D_i, D'_i . Since w is a curve in $\mathcal{R}$, hence we have,

$$\inf_{\sigma \in \Phi_i} \int_{\sigma} |\rho_{\epsilon, w}| \geq \inf_{v \in V_i} \int_v |\rho_{\epsilon, w}|$$

Since for $v \subset \mathcal{R}$ we have $\int_v |\rho_{\epsilon, w}| < \int_v |\rho_{[c]}| - \delta'_{\epsilon}$ for sufficiently small ϵ . This implies,

$$\begin{aligned} \mathcal{E}_{\mathcal{R}_i}(\Phi_i) &> \frac{\inf_{v \in V_i} \left(\int_v |\rho_{[c]}| - \delta'_{\epsilon} \right)^2}{\left(\int_{C_i \cup C'_i} |\rho_{[c]}| + \delta_{\epsilon} \right)^2} \geq \frac{\inf_{v \in V_i} \left(\int_v |\rho_{[c]}| - \delta'_{\epsilon} \right)^2}{\left(2 \int_{C_i} |\rho_{[c]}| + \delta_{\epsilon} \right)^2} \\ &> \frac{\inf_{\beta \in D(\alpha_i)} \ell^2(\beta) - \delta''_{\epsilon}}{4\ell^2([c]_i) + \delta'''_{\epsilon}}. \end{aligned}$$

Since $\delta''_{\epsilon}, \delta'''_{\epsilon}$ can be made arbitrarily small by choose ϵ sufficiently small, the last inequality implies,

$$\mathcal{E}_{\mathcal{R}_i}(\Phi_i) \geq \frac{1}{4} Q([c]_i).$$

Let g be the Mobius transformation so that $g\gamma_i g^{-1}$ of fixed points $0, \infty$. We have $g(\mathcal{R}_i) = A_i$ is annulus centered at origin of radii $r_2 > r_1$. Since,

$$g^* \mathcal{E}_{\mathcal{R}_i}(\Phi_i) = \mathcal{E}_{A_i}(g(\Phi_i)) = \frac{1}{2\pi} \log\left(\frac{r_2}{r_1}\right).$$

Also note that the translation length of $g\gamma_i g$ is $\log(\frac{r_2}{r_1})$, i.e: $T_{[c]_i} = \log(\frac{r_2}{r_1})$. Hence by conformal invariance of $\mathcal{E}_{\mathcal{R}_i}(\Phi_i)$ we have, $T_{[c]_i} \geq \frac{\pi}{2}Q([c]_i)$. $\square$

6 Pants decomposition and bound of $Q([c])$

Next we show the existence of homological basis which gives some lower bounds for the Rational Norm $Q([c])$.

Lemma 6.1. *There exists $[c] \in BH_1$ such that*

$$Q([c]) > \frac{2\lambda_g}{\pi}g \log(g),$$

for some $\lambda_g > 2$, when $g = 2$ and $\lambda_g > 3$ if $g > 2$.

Proof. We will choose $[c]$ which is of relatively short length to it's β_i^* by compare arcs on pair of pants. The case $g = 1$ is trivial. Assume $g = 2$. For $c \in [c]$, denote by $c = \{\alpha_1, \alpha_2\}$ with α_i the non-separating curves and it's dual curves (intersection $\langle \alpha_i, \beta_j \rangle = \delta_{ij}$) by β_i .

Since $g = 2$, take $\{\alpha_1, \alpha_2, \beta_1, \beta_2\}$ then, $\frac{\ell(\alpha_i)}{\ell(\beta_i)} < 1$ or inverse is < 1 for $1 \leq i \leq 2$. So we can always choose c so that $\frac{\ell(\beta_i)}{\ell(\alpha_i)} > 1$ for $1 \leq i \leq 2$. Hence we have $\sum Q([c]_i) > 2$ In addition, we have some $\lambda_g > 2$.

For $g > 2$, we need to decompose R_g into $2g - 2$ pair of pants and estimate $\ell(\beta)/\ell(\alpha)$ on pant components.

Given $[c] \in BH_1$ we complete $[c]$ with separating curves and let $P = \{P^k\}, 1 \leq k \leq 2g - 2$ denote the associated pants decomposition. For each $P^k \in P$ we cut into two hexagonal pieces and we mark it by border geodesic arcs by $B_k = \{g_k, b_k, c_k, c'_k, e_k, f_k, g'_k\}$. The arcs are: a_k right geodesic arc; b_k top connector geodesic arc; c_k top left geodesic arc, d_k top right left geodesic arc; e_k middle connector geodesic arc; f_k bottom geodesic arc; g_k left geodesic arc. Conversely, given any $2g - 2$ pair of pants decomposition of R_g there exists homological basis $\{\alpha_i, \beta_i\}, 1 \leq i \leq g$ which are not separating curves. To compare homological length we use idea of elementary arcs in [7]. A elementary arc e is a arc on P^k with end points lie on the boundary of P^k such that it intersect border geodesic arcs B_k at most two points in the interior of P^k . By definition all border geodesic arcs in B_k are elementary arcs.

The homological curve that we will be looking for must have minimal, although not necessarily zero Dehn twist, since by triangle inequality one can always shorten a curve by reduce its twist. We want to be able to show that there exists $[c]$ such that α_i , the cut curves can be made successively relatively short compare to its, dual β_i curves.

The idea is that, if there are no such $[c]$ exists on R_g then we will have an contradiction with the hyperbolicity of R_g . This contradiction is reached through length computations of arcs on P^k . To do so, we will compute relative $\frac{\ell(e)}{\ell(e')}$ for different pair of e, e' and show that it under conditions have sufficient bounds on them. These bounds will allow us to construct curves α_i which must satisfies Lemma 6.1. Of course to bound Q , one must keep in mind that our curve have minimal Dehn twist, otherwise this relative length can be made arbitrarily large or small but won't provide any meaningful bounds of Q .

Denote $a_k \in \{g_k, g'_k\}, d_k \in \{c_k, c'_k\}$ for $1 \leq k \leq 2g - 2$. We make the convention that given a_k and d_k we set a'_k, d'_k to be the other arc in the pair collection.

Assume that the lemma is false. There are several cases that we need to consider. We first show that there must exists a P of R_g such that there exists $P^k \in P$ with $\frac{\ell(d_k)}{\ell(b_k)} > \log^{4/5}(2g)$:

Lemma 6.2. *Assume Lemma 6.1 is not true. For every P there must exists some $P^k \in P$ such that, $\frac{\ell(d_k)}{\ell(b_k)} \geq \log^{4/5}(2g)$.*

Proof. We prove by contradiction. So we assume all P pants decompositions of R_g have a P_k such that $\frac{\ell(d_k)}{\ell(b_k)} < \log^{4/5}(2g)$.

Let $\kappa > 1$. Consider the following two cases:

- (A) : $\ell(a_k)\ell(b_k) \geq \kappa \log(2g)$, for all k .
- (B) : There exists some k such that $\ell(a_k)\ell(b_k) < \kappa \log(2g)$.

Case (A) : Cut P_k into hexagon and it follows from hyperbolic hexagonal [10] formulas we have, $\sinh(\ell(e_k)) \sinh(\ell(b_k)/2) = \cosh(\ell(a_k))$. This gives:

$$\ell(e_k) \leq \ell(a_k) + \sinh^{-1}\left(\frac{3}{\ell(b_k)}\right)$$

which implies,

$$\frac{\ell(a_k)}{\ell(e_k)} \geq \frac{\ell(a_k)}{\ell(a_k) + \sinh^{-1}\left(\frac{1}{\ell(b_k)}\right)}.$$

Since $\ell(a_k)\ell(b_k) \geq \kappa \log(2g)$, and $g \geq 3$ we have from the above inequality,

$$\frac{\ell(a_k)}{\ell(e_k)} \geq \frac{\ell(a_k)}{\ell(a_k) + \sinh^{-1}(\frac{1}{\ell(b_k)})} > f(\kappa).$$

Note $f(\kappa)$ is increasing function of κ and $f(\kappa) < 1$.

Let $e_{\hat{k}}$ be such that $\ell(e_{\hat{k}}) = \min_{1 \leq k \leq 2g-2} \{\ell(e_k)\}$. If $\hat{k} \leq g-1$ then we set $\alpha_1 = d_1 \cup d_{1'}$. Here $d_{k'}$ is the cut image curve of d_k , so $\ell(d_k) = \ell(d_{k'})$. On the other hand, if $\hat{k} > g-1$ then, we set $\alpha_1 = d_{2g-2} \cup d_{2g-2'}$. In either case we set, $\alpha_i = 2e_{i-1} \cup 2e_i$ for $i \geq 2$.

It follows from our choice of α_i we have, the curve homotopic to β_i must have arcs homotopic to a_i arcs. Now if $\hat{k} \leq g-1$ then set $\bar{\beta}_1 = \cup_{2g-2 \geq i \geq \hat{k}} (a_i \cup a_{i'})$, and if $\hat{k} > g-1$ then we set $\bar{\beta}_1 = \cup_{1 \leq i \leq \hat{k}} (a_i \cup a_{i'})$. And for $i \geq 2$ we also set $\bar{\beta}_i = \cup_{2g-2 \geq i} (a_i \cup a_{i'})$. Then it follows that we must have,

$$\frac{\ell(\beta_1)}{\ell(\alpha_1)} \geq \frac{\ell(\bar{\beta}_1)}{\ell(\alpha_1)} \geq \sum_{1 \leq i \leq g-1} f(\kappa) = f(\kappa)(g-1).$$

Hence we have,

$$\sum_{1 \leq i \leq 2g-2} \frac{\ell^2(\beta_i)}{\ell^2(\alpha_i)} > (f(\kappa)(g-1))^2, \quad \text{for } g \geq 3.$$

Since $\lim_{\kappa \rightarrow \infty} f(\kappa) = 1$ is increasing function so, there exists κ_o such that for $\kappa \geq \kappa_o$ we have for $g = 3$, $(f(\kappa)(2))^2 > \frac{3\pi}{2} \log(6)$. Now by the fact that,

$$\frac{\pi (f(\kappa)(g-1))^2}{2g \log(2g)}, \quad \text{is increasing function of } g.$$

Hence we have $Q([c]) > \frac{2}{\pi} g \log(2g)$ for $[c]$ consists of the chosen curves, which is a contradiction.

Next we consider case **(B)** : $\frac{\kappa \log(2g)}{\ell(a_{k_m})} > \ell(b_{k_m}) > \frac{\ell(d_{k_m})}{\log^{4/5}(2g)}$ for some k_m .

Now if $\ell(a_{k_m}) \geq \kappa \log(2g)$ for all the k_m then,

$$\frac{\ell(a_{k_m})}{\ell(d_{k_m})} > \frac{\ell(a_{k_m})}{\log^{4/5}(2g)} > 1.$$

Let $\hat{k}_m$ such that $\ell(d_{\hat{k}_m}) = \min_{k_m} \ell(d_{k_m})$. Here we choose as following:

Let $|\{k_m\}|$ denote number of elements of the collection. If $|\{k_m\}| \geq g-1$ then we choose,

$$\alpha_1 = d_{\hat{k}_m} \cup d_{\bar{k}_m}, \bar{\beta}_1 = \cup_{j \leq 2g-2} a_j \cup a_{j'},$$

and $\alpha_j = 2e_j \cup 2e_{j+1}$ for $j \neq k$.

From our choice of α_i , the curve β_1 must have arcs homotopic to a_j curves. Hence $\ell(\beta_1) \geq \ell(\bar{\beta}_1)$ and we have,

$$\frac{\ell(\beta_1)}{\ell(\alpha_1)} \geq \frac{\ell(\bar{\beta}_1)}{\ell(\alpha_1)} > \sum_{1 \leq j \leq 2g-2} 1 = 2g-2.$$

Hence for this basis we have, $\sum_{1 \leq i \leq g} Q([c]_i) > (2g-2)^2$. Since

$$\frac{(2g-2)^2}{g \log(2g)} > 1, \quad g \geq 3,$$

we have contradiction.

On the other hand if $|\{k_m\}| < g-1$ then we choose in combination with case (A): Let $\{d_{\hat{k}}\}$ be elements of $\{d_{k_m}\}$ such that among all curves β_i defined in (A) consist of $a_k \cup a_{k'}$ which do not intersect $d_{\hat{k}}$, gives $\frac{\ell(\beta_{\hat{k}})}{\ell(\alpha_{\hat{k}})}$ maximal value. Let d_{k^*} be the minimal length curve of $d_{\hat{k}}$. Then we have,

$$\begin{aligned} \sum_{1 \leq i \leq g} \frac{\ell^2(\beta_i)}{\ell^2(\alpha_i)} &\geq \frac{\ell^2(\bar{\beta}_{\hat{k}})}{\ell^2(\alpha_{\hat{k}})} + \frac{\ell^2(\bar{\beta}_{k^*})}{\ell^2(\alpha_{\bar{k}})} \\ &> \left(f(\kappa)(g - \frac{|\{k_m\}|}{2} - 1) \right)^2 + \left(\frac{|\{k_m\}|}{2} - 1 \right)^2 \end{aligned}$$

Hence by previous estimates we have $Q([c])$ satisfies the inequality, which gives contradiction.

Now suppose that $\ell(a_{k_m}) < \kappa \log(2g)$ for some of the m . We consider this as the Case (C). Here we breakdown the case (C) into two subcases:

- (C₁) : $\frac{\ell(a_k)}{\ell(b_k)} \geq 1$, for all k
- (C₂) : $\frac{\ell(a_k)}{\ell(b_k)} < 1$, some k_n .

Consider $(\mathbf{C}_1)$: In this case we have by our global condition $\ell(b_k) > \frac{\ell(d_k)}{\log^{4/5}(2g)}$ implies, $\frac{\ell(a_k)}{\ell(d_k)} > \frac{1}{\log^{4/5}(2g)}$.

We set $\bar{\beta}_1 = \cup_{2g-2 \geq j} a_j \cup a_{j'}$, $\alpha_1 = d_{\hat{k}} \cup d_{\hat{k}}$, where $\ell(d_{\hat{k}}) = \min_k \{\ell(d_k)\}$. And we set $\alpha_i = 2e_{i-1} \cup 2e_i$ for $i \geq 2$. Choose $\beta_{i \geq 2}$ so that $\{\alpha_i, \beta_i\}_{1 \leq i \leq g}$ form a basis. Similarly as before we have,

$$\frac{\ell(\beta_1)}{\ell(\alpha_1)} \geq \frac{\ell(\bar{\beta}_1)}{\ell(\alpha_1)} > \frac{2g-2}{\log^{4/5}(2g)}.$$

Since,

$$\frac{\pi(2g-2)^2}{2g \log^{14/5}(2g)} > 1, \quad \text{for } g \geq 2$$

and it is increasing function of g , it follows that we have $Q([c])$ satisfies the inequality which give us a contradiction.

Consider $(\mathbf{C}_2)$. In this case we have, $\ell(a_{k_n}) < \ell(b_{k_n})$.

By the hyperbolic identity $\sinh(\ell(e_k)) \sinh(\frac{\ell(b_k)}{2}) = \cosh(\ell(a_k))$ we have,

$$\frac{\ell(a_{k_n})}{\ell(e_{k_n})} > \frac{\ell(a_{k_n})}{\sinh^{-1} \left(\frac{\cosh(\ell(a_{k_n}))}{\sinh(\frac{\ell(a_{k_n})}{2})} \right)}.$$

Here we further subdivide into cases as:

- $(\mathbf{C}'_2) : \ell(a_{k_n}) \geq \frac{\rho}{\sqrt[3]{2g}}$.
- $(\mathbf{C}''_2) : \ell(a_{k_n}) < \frac{\rho}{\sqrt[3]{2g}}$.

For $(\mathbf{C}'_2)$, we have from the above inequality after some simple computations we have, $\frac{\ell(a_{k_n})}{\ell(e_{k_n})} > \left(\frac{\rho}{\sqrt[3]{2g}} \right)^{\frac{11}{10}}$. Note that this inequality holds for $\ell(a_{k_n}) \geq \frac{\rho}{\sqrt[3]{2g}}$ and $g \geq 3$.

Set $\alpha_1 = d_{\hat{k}} \cup d_{k'}$, $\bar{\beta}_1 = \cup_{1 \leq j \leq 2g-2} a_j$ and $\alpha_i = e_{2i-1} \cup e_{2i}$, $\bar{\beta}_i = \cup_{i \leq j \leq 2g-2} a_j \cup a_{j'}$ and $\bar{\beta}_{i-1} = \cup_{j \leq i-1} a_j \cup a_{j'}$, for $i \neq \hat{k}$. Then by similar computations as above then show that this basis $[c]$ satisfies,

$$\sum_{1 \leq i \leq g} \frac{\ell^2(\bar{\beta}_i)}{\ell^2(\alpha_i)} \geq \frac{4(g-|n|-1)^2}{\log^2(2g)} + \frac{\rho^{\frac{22}{10}} |n|^2}{(2g)^{11/30}}.$$

This inequality follows from that the maximal number of a_{k_n} arcs that don't pass through $d_{\hat{k}}$ is $|n|/2$. Here $|n|$ is the number of k_n .

This implies that, if $|n| > \frac{g}{2}$ then, we can always choose curves α_i as above so that, $\sum_{1 \leq i \leq g} \frac{\ell^2(\beta_i)}{\ell^2(\alpha_i)} > \frac{\rho^{\frac{22}{10}} g^{49/30}}{2^{71/30}}$. By setting $\rho = \frac{6}{10}$ we have,

$$\left(\pi \frac{\rho^{\frac{22}{10}} g^{49/30}}{2^{71/30}}\right) / (2g \log(2g)) > 1, \quad \text{for } g \geq 3.$$

It follows that $Q([c]) >$ gives our contradiction.

Case (**C**₂''). Here we have, $\ell(a_{k_n}) < \frac{\rho}{\sqrt[3]{2g}}$. Note by the global condition $\ell(d_k) < \ell(b_k) \log^{4/5}(2g)$ we also have, $\ell(b_{k_n}) < \frac{\kappa \log(2g)}{\ell(a_{k_n})}$.

we have by hexagonal hyperbolic formula,

$$\cosh(\ell(a_{k_n})) = \sinh(\ell(a_{k_n})) \sinh(b_{k_n}) \cosh(\ell(d_{k_n})) - \cosh(\ell(a_{k_n})) \cosh(\ell(b_{k_n}))$$

gives

$$\ell(d_{k_n}) = \cosh^{-1} \left(\frac{\cosh(\ell(a_{k_n}))(1 + \cosh(\ell(b_{k_n})))}{\sinh(\ell(a_{k_n})) \sinh(\ell(b_{k_n}))} \right).$$

by our conditions this implies,

$$\begin{aligned} \ell(d_{k_n}) &\geq \cosh^{-1} \left(\frac{\cosh(\ell(a_{k_n}))(1 + \cosh(\frac{\kappa \log(2g)}{\ell(a_{k_n})}))}{\sinh(\ell(a_{k_n})) \sinh(\frac{\log(2g)}{\kappa \ell(a_{k_n})})} \right) \\ &\geq \cosh^{-1} \left(\frac{\cosh(\frac{6}{\sqrt[3]{2g}})(1 + \cosh(\frac{\kappa \log(2g)}{\ell(a_{k_n})}))}{\sinh(\frac{6}{\sqrt[3]{2g}}) \sinh(\frac{\kappa \log(2g)}{\ell(a_{k_n})})} \right) \\ &> \frac{17}{10} \log(2g) \end{aligned}$$

By $\ell(d_k) < \ell(b_k) \log^{4/5}(2g)$ implies that, $\ell(b_{k_n}) > \frac{17}{10} \log^{1/5}(2g)$. We have,

$$\frac{\ell(b_{k_n})}{\ell(e_{k_n})} > \frac{(2) \frac{17}{10} \log^{1/5}(2g)}{\sinh^{-1} \left(\frac{\cosh(\frac{6}{\sqrt[3]{2g}})}{\sinh(\frac{17}{10} \log^{1/5}(2g))} \right)}$$

With some computations one shows that, $\frac{\ell(b_{k_n})}{\ell(e_{k_n})} > 3 \log(2g)$.

Here we set $\alpha_1 = \cup_{1 \leq j \leq 2g-2} a_j$ and $\alpha_i = 2e_{2i-1} \cup 2e_{2i}$ for all the i that within the collection of P_{k_n} . We select the remaining curves α_j which are not consist of e_j . By construction it follows that, β_i the geodesic curve must have arcs homotopic to a_i and, $\sum_{i \leq i \leq g} \frac{\ell(\beta_i)}{\ell(\alpha_i)} > \frac{9|n|}{4} \log^2(2g)$. Now if $|n| \geq \frac{g}{2}$ then we have, $Q([c]) > \frac{2}{\pi} g \log(2g)$. On the other hand, if $|n| < \frac{g}{2}$ we can then choose curves given by previous construction. Hence we always curves that gives our contradiction, this completes our prove of the result. $\square$

Now we establish the induction process of showing:

Corollary 6.3. *Assume Lemma 6.1 is false. Let $1 < i \leq g$. Suppose there exists P^{k_j} for every $j < i$ such that, $\frac{\ell(d_{k_j})}{\ell(b_{k_j})} \geq \log^{4/5}(2g - 2j + 2)$. Then there must exists $P^{k_i} \in P - \cup_{j < i} P^{k_j}$ such that $\frac{\ell(d_{k_i})}{\ell(b_{k_i})} \geq \log^{4/5}(2g - 2i + 2)$.*

Proof. The pants $P^{k_i} \in P - \cup_{j < i} P^{k_j}$ consists of decomposition of surface of genus $g - i + 1$. It follows from Lemma 6.2, there exists d_{k_i}, b_{k_i} such that $\frac{\ell(d_{k_i})}{\ell(b_{k_i})} \geq \log^{4/5}(2(g - i + 1))$. $\square$

Now we can finish the proof of Lemma 6.1: If the Lemma is false then, we choose the collection of geodesic curves on surface R_g represented by $\alpha_i = b_i \cup b_{i'}$ for $1 \leq i \leq g$. Since the shortest geodesic β_i intersecting α_i is given by either $c_i \cup c'_i$ or $c'_i \cup c_{i'}$, i.e. $\beta_i = d_i \cup d_{i'}$. Hence it follows from Corollary 6.3 we have that,

$$\frac{\ell(\beta_i)}{\ell(\alpha_i)} \geq \log^{4/5}(2g - 2i + 2), \quad \text{for } 1 \leq i \leq g.$$

After simple computation one shows that, $\sum_{i=1}^g \log^{8/5}(2g - 2i + 2) > \frac{2}{\pi} g \log(2g)$. Hence $Q([c]) > \frac{4}{\pi} g \log(g)$, a contradiction. $\square$

Proposition 6.4. *There exists a $[c] \in BH_1$ such that,*

$$Q(\theta.[c]) > \frac{2\lambda_g}{\pi} g \log(g), \quad \text{for all } \theta \in \text{stab}(\phi).$$

Remark 6.5. Note that under the action of θ we could have i permuted, hence only the summation of the above inequality is preserved. We call any such $[c]$ satisfies the inequality, *positive*. Call $[c]$ *invariantly positive* if having property of Lemma 6.4.

Proof. We need to examine θ action on elementary curves on pants decomposition. Note that it is enough to show it is true for generators of $stab(\phi)$. We can written elements of $SP(2g, \mathbb{Z})$ as composition of several types of elementary symplectic matrices. For elements of $stab(\phi)$, which is subgroup generated by elementary matrices which do not intertwiners the α_i and β_j basis. These elementary matrix correspond to a Nielsen transformation on the generators of $\phi([c])$.

The idea is that, take $[c]$ positive which exists by Lemma 6.1, by similar computations as in the proof of Lemma 6.1 using elementary arcs, if $\theta.[c]$ is not positive then, we can appropriately modify the original curve of $[c]$ to a new $[c']$ so that this, $[c']$ will only increase Q under θ . This is achieved by compare different elementary arc length as we have done previously.

Let E_{ij} denote a elementary matrix such that E_{ij} map $[c]$ into basis with α_i replaced by $\alpha_i + \alpha_j$ and β_j replaced by $\beta_j - \beta_i$, and rest unchanged. Example E_{12} for $g = 3$ of $SP(6, \mathbb{Z})$:

$$E_{12} = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We need to consider cases on pair pants. Assume the statement is false and there exists l, m such that $\sum_j Q[E_{lm}.[c]_j] < \frac{2}{\pi}g \log(2g)$ for all positive $[c]$. It follows from the proof of Lemma 6.1, it is suffice to consider α_l, α_m are curves consist of elementary arcs. Let h_{lm} denote the geodesic curve representation of homological class $[\alpha_l + \alpha_m]$. Let $[f_{lm}]$ be a class of curves which is canonical dual to $[h_{lm}]$ and non-intersecting to rest of basis. We denote the f_{lm} to be the geodesic representative of $[f_{lm}]$.

Take a $[c]$ provided by Lemma 6.1. We will show that either this $[c]$ is invariantly positive or it implies another $[\hat{c}]$ is invariantly positive. To do so we will examine all possible cases of $[c]$. We continue to use our notations and definitions given in the proof of Lemma 6.1. We need to examine all possible curves given by the $E_{lm}.[c]$.

As in the Lemma 6.1, first assume that we have $\frac{\ell(d_k)}{\ell(b_k)} < \log^{4/5}(2g)$ for all P^k . Note that we have some positive $[c]$. Since h is geodesic closed curve homotopically connects b_l, b_m , the length is bounded above by sum of these

curves and twice the connecting geodesic curve. As in the proof of Lemma 6.1, if $\ell(a_k)\ell(b_k) \geq \log(2g)$ for all k , then $\ell(e_l) + \ell(e_m) > \ell(d_l) + \ell(d_m)$. So if we set $\delta = \ell(h_{lm}) - (\ell(b_l) + \ell(b_m))$, then $\ell(f_{lm}) \geq \ell(e_l) + \ell(e_m) + \delta$. Hence we have, $\frac{\ell(f_{lm})}{\ell(h_{lm})} \geq \frac{g}{\log(2g)}$, and we have $E_{lm} \cdot [c]$ positive. On the other hand if, $\ell(a_k)\ell(b_k) < \log(2g)$ for some P^k contains b_l, b_m then, we have a curve α which consists of a_k as elementary arcs and $\frac{\ell(d_j)}{\ell(\alpha)} > \frac{g}{\log(2g)}, j \in \{l, m\}$. We replace one of the original α_l with α to form basis $[\hat{c}]$. It is obvious that $[\hat{c}]$ is positive.

For simplicity we use same notation for the geodesic representation curves for $[\hat{c}]$. Then h have α_l a Dehn twist around b_m which trace off from a_k arcs. The geodesic f_{lm} consists of arcs homotopic to arcs of β_m, β_l, b_l . Hence $\ell(f_{lm}) \geq \frac{1}{2}(\ell(\beta_m) + \ell(\beta_l)) + \ell(b_l)$. This implies we have lower bound of

$$\frac{\ell(f_{lm})}{\ell(b_l)} \geq \frac{\ell(\beta_m)}{\ell(b_l)} + 1.$$

Hence we have, $E_{lm} \cdot [\hat{c}]$ positive. By repeat this replacement for rest of curves of $[c]$ if necessary we can claim that $E_{lm} \cdot [\hat{c}]; 1 \leq l, m \leq g$. satisfies same inequality. Note that $E_{lm} \cdot E_{l'm'} \cdot [\hat{c}]$ will increase Dehn twist which will increase the lower bound by twist number. i.e. denote $f_{l_1 m_1 \dots l_j m_j}$ be the geodesic representative of $[f]$ corresponds to $E_{l_1 m_1} \dots E_{l_j m_j} \cdot [\hat{c}]$ we have,

$$\frac{\ell(f_{l_1 m_1 \dots l_j m_j})}{\ell(b_l)} \geq \frac{\ell(\beta_m)}{\ell(b_l)} + j.$$

Hence we have, $[\hat{c}]$ is invariantly positive.

Now it follows from Corollary 6.3 and Lemma 6.2 that, every pants decompositions have cut curves α_i such that, β_i have the property of $\frac{\ell(\beta_i)}{\ell(\alpha_i)} \geq \log^{4/5}(2g), 1 \leq i \leq g$. This implies that there exists a invariantly positive $[c]$. $\square$

From Lemma 6.1 and Lemma 6.4 we can prove the following inequality of the Schottky length,

Proposition 6.6. *There exists $[c] \in BH_1$ such that,*

$$\mathcal{T}_{[c]} > \lambda_g \log(g).$$

Proof. Let $[c]$ be given by Lemma 6.1. By Lemma 5.5 we have,

$$T_{[c]_i} \geq \frac{\pi}{2} Q([c]_i).$$

Since $\sum_i Q([c]_i) > \frac{2\lambda_g}{\pi} g \log(g)$, this implies that $\sum_{i=1}^g T_{[c]_i} > \lambda_g g \log(g)$. Since $\mathcal{T}_{[c]} = \frac{1}{g} \sum_{i=1}^g T_{[c]_i}$, hence the above strict inequality implies the result. $\square$

7 Proof of Theorem 1.1

Let $\llbracket c \rrbracket \in \hat{B}H_1$. For $[c] \in \llbracket c \rrbracket$, $\omega_{[c]}$ is an generators set of $\Gamma_{[c]}$. Since $\Gamma_{[c]} = \phi([c]), \forall [c] \in \llbracket c \rrbracket$ and $\bigcup_{[c] \in \llbracket c \rrbracket} \omega_{[c]} = \mathbb{W}_\Gamma$ we have our next corollary,

Proposition 7.1. *Let $[c]$ be given by Proposition 6.6. Then $\|\mathbb{W}_\Gamma\|_x > \lambda_g \log(g)$.*

Proof. It follows from Proposition 6.6 and $\mathfrak{w}_x(\omega_{[c]}) = \mathcal{T}_{[c]}$, we have:

$$\mathfrak{w}_x(\omega_{[c]}) > \lambda_g \log(g).$$

Since $\bigcup_{[c] \in \llbracket c \rrbracket} \omega_{[c]} = \mathbb{W}_\Gamma$, by the invariant positivity result of Lemma 6.4 we have :

$$\mathfrak{w}_x(\omega_{[c]}) > \lambda_g \log(g), \quad \forall [c] \in \llbracket c \rrbracket.$$

Hence we have:

$$\|\mathbb{W}_\Gamma\|_x = \inf_{\omega_{[c]} \in \mathbb{W}_\Gamma} \mathfrak{w}_x(\omega_{[c]}) > \lambda_g \log(g).$$

$\square$

Proof. Theorem 1.1:

For a R_g , it follows from Proposition 7.1, there exists homological marking $[c] \in BH_1$ of R_g such that, $\Gamma_{[c]} = \phi([c])$, the covering Schottky group satisfies, $\|\mathbb{W}_\Gamma\|_x > \lambda_g \log(g)$ for all $\mathbb{H}^3$. Hence by Corollary 4.5 and Corollary 4.6, and note that for such $[c]$ we have $\lambda < 2$, which implies $\mathfrak{D}_{\Gamma_{[c]}} < 1$.

$\square$

Finally, we give two obvious applications of our theorem. The first application Corollary 7.2, address a folklore question that was originally due to Bers.

Corollary 7.2. *If $H_c \geq 1$ then, for all $[R_g] \in \mathcal{M}_g$, $\pi_S^{-1}([R_g])$, have classical fiber.*

Proof. Assume that $H_c \geq 1$, so $\tau_c \leq 1$. Since $\tau_c \leq 1$, Proposition 3.2 and Theorem 2.1 implies that Γ is classical Schottky group. Hence for $[R_g] \in \mathcal{M}_g$, $\pi_S^{-1}([R_g]) \cap \mathfrak{J}_{g,o} \neq \emptyset$. Therefore, $\pi_S|_{\mathfrak{J}_{g,o}}$ is surjective. $\square$

The second application is presentation of period matrix of R_g in Schottky coordinates. It's well known theorem of Torelli that there exists a injective map from $\mathcal{M}_g$ into Siegel's space of symmetric $g \times g$ matrix over $\mathbb{C}$. This is given by the P_{mn} period matrix of R_g . Many beautiful theorems has been proved on P_{mn} , such as Buser-Sarnak's theorem [6] states that the locus of Jacobians lie in very small neighborhood of the boundary of space of principally polarized abelian varieties for large genus R_g .

It's well known fact [4], that P_{mn} can be represented in local coordinates of $\mathfrak{J}_g$. Recall that local coordinates of $\mathfrak{J}_g$, which is $3g-3$ -dimensional complex manifold, are given by variables $\lambda_i, z_{-,i}, z_{+,i}, 1 \leq i \leq g-1$ multiplier and two fixed points respectively. Recall, given $z_1, z_2, z_3, z_4 \in \mathbb{C}$, the cross ratio is: $[z_1, z_2, z_3, z_4] = \frac{(z_1-z_3)(z_2-z_4)}{(z_1-z_4)(z_2-z_3)}$. Let $\Gamma \in \mathfrak{J}_g$ be generated by $\langle \gamma_1, \dots, \gamma_g \rangle$, we denote by Γ_l the subgroup of Γ generated by γ_l . Then the following *formal* presentation of P_{mn} is well known and go back to Schottky himself [4, 21]:

$$P_{mn} = \sum_{\gamma \in \Gamma_n \setminus \Gamma / \Gamma_m} \log[z_{-,n}, \gamma z_{-,m}, z_{+,n}, \gamma z_{+,m}] + \delta_{mn} \log \lambda_n.$$

However, even though P_{mn} has been formally known for a very long time but, in general as P_{mn} is infinite sum, so it is not always convergent for a arbitrary $\Gamma \in \mathfrak{J}_g$, and it is not known in general. In this respect, we have our second simple application of Theorem 1.1:

Corollary 7.3. *Let $[R_g] \in \mathcal{M}_g$. There exists $\Gamma \in \mathfrak{J}_g$ such that the period matrix of P_{mn} of $[R_g]$ is given by the above presentation.*

Proof. The only thing needs to be verified is that P_{mn} is convergent for some $\Gamma \in \pi_S^{-1}([R_g])$. By Theorem 2.1, we have a $\hat{\Gamma} \in \pi_S^{-1}([R_g])$ such that $\mathfrak{D}_{\hat{\Gamma}} < 1$.

Note that P_{mn} is obtained as integral around canonical basis cycles of $H_1(R_g, \mathbb{Z})$ of $\hat{\Gamma}$ -invariant holomorphic cocycles on $\Omega_{\hat{\Gamma}}$:

$$\omega_n(z) = \sum_{\gamma \in \hat{\Gamma}_n \setminus \hat{\Gamma}} d \log \left(\frac{z - \gamma z_{+,n}}{z - \gamma z_{-,n}} \right).$$

It follows from the residue formula we have P_{mn} .

By change of variable we have ω_n is convergent if the Poincare series: $\sum_{\gamma \in \hat{\Gamma}_n \setminus \hat{\Gamma}} |\gamma'(z)|$ is convergent. Since $\mathfrak{D}_{\hat{\Gamma}} < 1$, we have $\sum_{\gamma \in \hat{\Gamma}_n \setminus \hat{\Gamma}} |\gamma'(z)|$ is convergent. Hence P_{mn} exists for $\hat{\Gamma}$. $\square$

E-mail: yonghou@princeton.edu

References

- [1] Agol, I. (2004). Tameness of hyperbolic 3-manifolds. Arxiv. <http://arxiv.org/abs/math/0405568>
- [2] Alvarez, V. Pestana, D. Rodriguez, J. (1999). Revista Math Ibero. Vol, 2. 353-427.
- [3] Ahlfors. L. (1960). Riemann surfaces. Princeton Press.
- [4] Baker, H. F. (1995). Abelian functions: Abel's theorem and the allied theory, including the theory of the theta functions. Cambridge Univ Press.
- [5] Bers, L. (1975). Automorphic forms for Schottky groups. Adv. Math. 16, 332-361.
- [6] Buser, P. Sarnak, P. (1994). On the period matrix of a Riemann surface of large genus. Invent. Math. Vol 117. pp. 27-56.
- [7] Buser, P. Seppala, M. (2002). Short homology bases and partitions of Riemann surfaces. Topology. 41, 863-871.
- [8] Coornaert. M and Papadopoulos, A. Symbolic dynamics and hyperbolic groups. Lecture Notes in Math. 1539, Springer.
- [9] Culler, M., Shalen, B.P. (1996). Free Kleinian groups and volumes of hyperbolic 3-manifolds. J. Diff. Geo., 44, 738-782.
- [10] A. Berdon. (1983). The Geometry of discrete groups, Springer-verlag.
- [11] Calegari, D. Gabai, D. (2004) Shrinkwrapping and the taming of hyperbolic 3-manifolds. Arxiv. <http://arxiv.org/abs/math/0407161>

- [12] Doyle, P. (1988) On the bass note of a Schottky group. *Acta Math.* 160, 249-284.
- [13] Farb, B. Margalit, D. (2011). *A Primer on mapping class groups*. Princeton Press.
- [14] Gromov, M. (1987). Hyperbolic Groups. In *essays in group theory*. Gersten (editor) M.S.R.I. Publ. 8. Springer, Berlin. pp 75-263.
- [15] Hejhal. D. (1975). On Schottky and Teichmuller spaces. *Adv. in. Math.* 15, pp133-156.
- [16] Hou, Y. (2014). All finitely generated Kleinian groups of small Hausdorff dimension is classical Schottky group. To appear *Adv. in. Math.* <http://arxiv.org/abs/1307.2677>
- [17] Hou, Y. (2001). Critical exponent and displacement of negatively curved free groups. *J. Diff. Geo.* 57, 173-195.
- [18] Hou, Y. (2010). Kleinian groups of small Hasudorff dimension are classical Schottky groups. I. *Geometry & Topology.* 14, 473-519.
- [19] Kapovich, M. (2000). *Hyperbolic Manifolds and Discrete Groups: Lecture on Thurston's Hyperbolization*, Birkhauser's series "Progress in mathematics".
- [20] Keen, L. etc.. (1995). *Lipa's Legacy*. *Contemp Math.* pp. 433.
- [21] Manin, Y. Drinfeld, G. (1973). Periods of p -adic Schottky groups. *J. Reine. Anger. Math.* 263, 239-247.
- [22] Marden, A. (1974). *Schottky groups and circles*. *Contributions to analysis* (a collection of papers dedicated to Lipman Bers). pp. 273-278. Academic Press.
- [23] Marden, A. (2007). *Outer circles: An introduction to hyperbolic 3-manifolds*. Cambridge Press.
- [24] Maskat. B. (1988). *Kleinian Groups*. *Grundlehren der math. Wiss.* 287. Springer.

- [25] McMullen, C. (1984). Complex Dynamics and Renormalization, *Annals of Math Studies.* 135. Princeton University Press.
- [26] McMullen, C. (1998). Hausdorff dimension and conformal dynamics, III: Computation of dimension. *American Journal of Mathematics.* Vol 120, No 4, 691-721.
- [27] Morgan, F. Johnson, D. (2000). Some sharp isoperimetric theorems for Riemannian manifolds. *Indiana Univ Math.*, Vol. 49. 1017-1041
- [28] Patterson, S.J. (1987). Measures on limit sets of Kleinian groups. *Analytical and Geometrical Aspects of hyperbolic space.* Cambridge Univ. Press. pp 291-323.
- [29] Phillips, R., Sarnak, P. The Laplacian for domains in hyperbolic space and limit sets of Kleinian groups. *Acta Math.*, 155:173-241
- [30] Sibner, R. (1965). Uniformization of symmetric Riemann surfaces by Schottky groups. *Trans.AMS.* 116.
- [31] Sullivan, D. (1982). Discrete conformal groups and measurable dynamics. *Bull. Amer. Math. Soc.* 6, 57-73.